

Ready to begin again with BIKTARVY®?

Pill is not actual size
(15 mm x 8 mm)



NOW FDA APPROVED
for people restarting treatment

Talk to your healthcare provider about
whether BIKTARVY could be right for you.

Whether you're starting, switching, or restarting treatment, BIKTARVY may be right for you.

BIKTARVY is a complete, 1-pill, once-a-day prescription medicine used to treat HIV-1 in adults and children who weigh at least 31 pounds. It can either be used in people who have never taken HIV-1 medicines before, or people who have received HIV-1 medicines in the past, or to replace their current HIV-1 medicines, and whose healthcare provider determines they meet certain requirements.

BIKTARVY does not cure HIV-1 or AIDS. HIV-1 is the virus that causes AIDS.

IMPORTANT SAFETY INFORMATION

What is the most important information I should know about BIKTARVY?


BIKTARVY may cause serious side effects:

- ▶ **Worsening of hepatitis B (HBV).** Your healthcare provider will test you for HBV. If you have both HIV-1 and HBV and stop taking BIKTARVY, your HBV may suddenly get worse. Do not stop taking BIKTARVY without first talking to your healthcare provider, as they will need to monitor your health, and may give you HBV medicine.

Who should not take BIKTARVY?

Do not take BIKTARVY if you take:

- ▶ dofetilide
- ▶ rifampin
- ▶ any other medicines to treat HIV-1



BIKTARVY®

bictegravir 50mg/emtricitabine 200mg/
tenofovir alafenamide 25mg tablets

Wherever you are in your journey, it may be possible to get back on treatment

No matter what got in the way of your HIV treatment, a conversation with your healthcare provider can guide you through next steps.

Have an open and honest conversation with your healthcare provider about why you may have stopped so you can work together to determine the best path forward for you.

Here are some topics to consider and discuss with your healthcare provider if they were part of the reason why you stopped:


- ◆ Side effects
- ◆ Mental health
- ◆ The size or number of pills
- ◆ Food requirements
- ◆ Keeping up with appointments
- ◆ Cost of the medicine
- ◆ Changes in your life situation



Talk to your healthcare provider about anything stopping you from staying on treatment so that you can work together to find the next best step for you.

Model portrayals





BIKTARVY[®]

bictegravir 50mg/emtricitabine 200mg/
tenofovir alafenamide 25mg tablets

Whether you're starting or
restarting treatment, BIKTARVY[®]
may be right for you



BIKTARVY is the **first and only, once-daily, single pill** to receive
an **additional FDA approval** for use in **people with HIV restarting
treatment**, when appropriate.

**Ask your healthcare provider if you can
restart HIV treatment with BIKTARVY.**



Not actual size
(15 mm x 8 mm)

BIKTARVY is:

- ♦ **A complete HIV-1 treatment**
- ♦ **1 small pill once a day**
- ♦ **Taken any time of day**
- ♦ **Taken with or without food**

BIKTARVY does not cure HIV-1 or AIDS. HIV-1 is the
virus that causes AIDS.

Person featured
is compensated
by Gilead.

Chad

Started on BIKTARVY
in 2018, still undetectable


IMPORTANT SAFETY INFORMATION (cont'd)

What are the other possible side effects of BIKTARVY?

Serious side effects of BIKTARVY may also include:

- ▶ **Changes in your immune system.** Your immune system may get stronger and begin to fight infections that may have been hidden in your body. Tell your healthcare provider if you have any new symptoms after you start taking BIKTARVY.
- ▶ **Kidney problems, including kidney failure.** Your healthcare provider should do blood and urine tests to check your kidneys. If you develop new or worse kidney problems, they may tell you to stop taking BIKTARVY.
- ▶ **Too much lactic acid in your blood (lactic acidosis),** which is a serious but rare medical emergency that can lead to death. Tell your healthcare provider right away if you get these symptoms: weakness or being more tired than usual, unusual muscle pain, being short of breath or fast breathing, stomach pain with nausea and vomiting, cold or blue hands and feet, feel dizzy or lightheaded, or a fast or abnormal heartbeat.

Please click to see Important Facts about [BIKTARVY](#), including an important warning on worsening of hepatitis B, on page 7.



BIKTARVY®

bictegravir 50mg/emtricitabine 200mg/
tenofovir alafenamide 25mg tablets

You don't have to go on this journey alone; members of your healthcare team can help

Whether you're just starting to think about getting back on treatment or you're ready to begin again, know that **there are many care team professionals out there who want to support you, such as:**

- ◆ Case managers
- ◆ Nurses
- ◆ Pharmacists
- ◆ Social workers
- ◆ Counselors
- ◆ Peer advocates

These professionals often work with many people living with HIV and are trained to help guide you through the process of re-entering HIV care. Talk to them openly about the reasons you may have stopped treatment.

When you're ready, they may be able to help you find a healthcare provider to see if restarting on BIKTARVY® is right for you.

Model portrayals



IMPORTANT SAFETY INFORMATION (cont'd)

What are the other possible side effects of BIKTARVY? (cont'd)

► **Severe liver problems**, which in rare cases can lead to death. Tell your healthcare provider right away if you get these symptoms: skin or the white part of your eyes turns yellow, dark "tea-colored" urine, light-colored stools, loss of appetite for several days or longer, nausea, or stomach-area pain.

The most common side effects of BIKTARVY in clinical studies were diarrhea (6%), nausea (6%), and headache (5%). Tell your healthcare provider if you have any side effects that bother you or don't go away.

What should I tell my healthcare provider before taking BIKTARVY?

► **All your health problems.** Be sure to tell your healthcare provider if you have or have had any kidney or liver problems, including hepatitis virus.

Please click to see Important Facts about [BIKTARVY](#), including an important warning on worsening of hepatitis B, on page 7.



Staying on track with treatment is important to your health

Taking your HIV treatment as prescribed is one of the most important things you can do to treat your HIV. If you're having concerns about taking your HIV medicines, talk openly with your healthcare provider and any other health professionals who support you.

Here are some ways to help you stay on treatment once you restart:



Make treatment part of your daily routine.

Be consistent about when you take your treatment and set a reminder in your phone. Ask your pharmacy to help you set up automatic refills for your prescription.



Join a support group. In addition to family and friends, there are HIV support groups. They can help people living with HIV share experiences and perspectives, reduce stigma, and manage feelings of isolation or depression.



Don't be afraid to ask for help. If something unexpected happens that makes it harder to keep up with treatment, reach out to your HIV care team. They want to help you overcome any obstacles to treatment.



Remember, you don't have to take this journey alone. It's possible to find the support and assistance you need to stay on treatment.


IMPORTANT SAFETY INFORMATION (cont'd)

What should I tell my healthcare provider before taking BIKTARVY? (cont'd)

- ▶ **All the medicines you take**, including prescription and over-the-counter medicines, antacids, laxatives, vitamins, and herbal supplements. BIKTARVY and other medicines may affect each other. Keep a list of all your medicines and show it to your healthcare provider and pharmacist, and ask if it is safe to take BIKTARVY with all of your other medicines.
- ▶ **If you are pregnant** or plan to become pregnant. Tell your healthcare provider if you become pregnant while taking BIKTARVY.
- ▶ **If you are breastfeeding** (nursing) or plan to breastfeed. Talk to your healthcare provider about the risks of breastfeeding during treatment with BIKTARVY.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088.

Please click to see Important Facts about [BIKTARVY](#), including an important warning on worsening of hepatitis B, on page 7.



BIKTARVY®

bictegravir 50mg/emtricitabine 200mg/
tenofovir alafenamide 25mg tablets

Don't let cost stand in your way

Insurance or no insurance, Gilead's Advancing Access® program may be able to help.

If you're eligible, Advancing Access® may be able to help you lower your co-pay to as little as \$0.* If you do not have insurance, you may be able to receive BIKTARVY® free of charge if you qualify for the Advancing Access Patient Assistance Program.



90% of people using the BIKTARVY co-pay program pay as little as \$0 a month.

*For commercially insured eligible patients only. Restrictions apply. Subject to change. See full terms and conditions at [GileadAdvancingAccess.com](https://www.gileadadvancingaccess.com).

Advancing Access specialists can help you understand your health insurance and Gilead medication costs. For more information, visit [GileadAdvancingAccess.com](https://www.gileadadvancingaccess.com) or call 1-800-226-2056, Monday-Friday, 9 AM to 8 PM ET.

Llame a la línea directa y pida hablar con un representante de habla hispana.



GILEAD

BIKTARVY, the BIKTARVY Logo, ADVANCING ACCESS, GILEAD, the GILEAD Logo, and GSI are trademarks of Gilead Sciences, Inc., or its related companies. © 2025 Gilead Sciences, Inc. All rights reserved. US-BVYC-0824 08/25

Pill is not actual size
(15 mm x 8 mm)



Find out how you can begin treatment again with BIKTARVY

When you're ready, talk to your healthcare provider about whether BIKTARVY may be the right treatment for you.

Visit [BIKTARVY.com/resources](https://www.biktarmvy.com/resources) for more information about BIKTARVY and resources to help support you on your restart journey.

Please click to see Important Facts about [BIKTARVY](#), including an important warning on worsening of hepatitis B, on page 7.



BIKTARVY®

bictegravir 50mg/emtricitabine 200mg/
tenofovir alafenamide 25mg tablets

(bik-TAR-vee)

MOST IMPORTANT INFORMATION ABOUT BIKTARVY

BIKTARVY may cause serious side effects, including:

- **Worsening of hepatitis B (HBV).** Your healthcare provider will test you for HBV. If you have both HIV-1 and HBV, your HBV may suddenly get worse if you stop taking BIKTARVY. Do not stop taking BIKTARVY without first talking to your healthcare provider, as they will need to check your health regularly for several months, and may give you HBV medicine.

ABOUT BIKTARVY

BIKTARVY is a complete, 1-pill, once-a-day prescription medicine used to treat HIV-1 in adults and children who weigh at least 31 pounds. It can either be used in people who have never taken HIV-1 medicines before, or people who have received HIV-1 medicines in the past, or to replace their current HIV-1 medicines, and whose healthcare provider determines they meet certain requirements.

BIKTARVY does not cure HIV-1 or AIDS. HIV-1 is the virus that causes AIDS.

Do NOT take BIKTARVY if you also take a medicine that contains:

- dofetilide
- rifampin
- any other medicines to treat HIV-1

BEFORE TAKING BIKTARVY

Tell your healthcare provider all your medical conditions, including if you:

- Have or have had any kidney or liver problems, including hepatitis.
- Are pregnant or plan to become pregnant.
- Are breastfeeding (nursing) or plan to breastfeed. Talk to your healthcare provider about the risks of breastfeeding during treatment with BIKTARVY.

Tell your healthcare provider about all the medicines you take:

- Keep a list that includes all prescription and over-the-counter medicines, antacids, laxatives, vitamins, and herbal supplements, and show it to your healthcare provider and pharmacist.
- Ask your healthcare provider or pharmacist about medicines that interact with BIKTARVY.

IMPORTANT FACTS

This is only a brief summary of important information about BIKTARVY® and does not replace talking to your healthcare provider about your condition and your treatment.

POSSIBLE SIDE EFFECTS OF BIKTARVY

BIKTARVY can cause serious side effects, including:

- Those in the “Most Important Information About BIKTARVY” section.
- Changes in your immune system.
- New or worse kidney problems, including kidney failure.
- Too much lactic acid in your blood (lactic acidosis), which is a serious but rare medical emergency that can lead to death. Tell your healthcare provider right away if you get these symptoms: weakness or being more tired than usual, unusual muscle pain, being short of breath or fast breathing, stomach pain with nausea and vomiting, cold or blue hands and feet, feel dizzy or lightheaded, or a fast or abnormal heartbeat.
- Severe liver problems, which in rare cases can lead to death. Tell your healthcare provider right away if you get these symptoms: skin or the white part of your eyes turns yellow, dark “tea-colored” urine, light-colored stools, loss of appetite for several days or longer, nausea, or stomach-area pain.
- **The most common side effects of BIKTARVY** in clinical studies were diarrhea (6%), nausea (6%), and headache (5%).

These are not all the possible side effects of BIKTARVY. Tell your healthcare provider right away if you have any new symptoms while taking BIKTARVY.

Your healthcare provider will need to do tests to monitor your health before and during treatment with BIKTARVY.

HOW TO TAKE BIKTARVY

Take BIKTARVY 1 time each day with or without food.

GET MORE INFORMATION

- This is only a brief summary of important information about BIKTARVY. Talk to your healthcare provider or pharmacist to learn more.
- Go to [BIKTARVY.com](https://www.biktarvy.com) or call 1-800-GILEAD-5.
- If you need help paying for your medicine, visit [BIKTARVY.com](https://www.biktarvy.com) for program information.



BIKTARVY, the BIKTARVY Logo, GILEAD, and the GILEAD Logo are trademarks of Gilead Sciences, Inc., or its related companies. Version date: July 2025
© 2025 Gilead Sciences, Inc. All rights reserved. US-BVYC-0824 08/25