

(des-KOH-vee)

## MOST IMPORTANT INFORMATION ABOUT DESCOVY

### DESCOVY may cause serious side effects, including:

- **Worsening of hepatitis B (HBV).** Your healthcare provider will test you for HBV. If you have both HIV-1 and HBV, your HBV may suddenly get worse if you stop taking DESCOVY. Do not stop taking DESCOVY without first talking to your healthcare provider, as they will need to check your health regularly for several months or give you HBV medicine.

## ABOUT DESCOVY

- DESCOVY is a prescription medicine that is used together with other HIV-1 medicines to treat HIV-1 in people who weigh at least 77 lbs (35 kg).
- **DESCOVY does not cure HIV-1 or AIDS.** HIV-1 is the virus that causes AIDS.

## BEFORE TAKING DESCOVY

### Tell your healthcare provider if you:

- Have or had any kidney or liver problems, including hepatitis.
- Have any other medical condition.
- Are pregnant or plan to become pregnant.
- Are breastfeeding (nursing) or plan to breastfeed. DESCOVY passes to your baby in your breast milk. Talk to your healthcare provider about the risks of breastfeeding during treatment with DESCOVY.

### Tell your healthcare provider about all the medicines you take:

- Keep a list that includes all prescription and over-the-counter medicines, vitamins, and herbal supplements, and show it to your healthcare provider and pharmacist.
- Ask your healthcare provider or pharmacist about medicines that should not be taken with DESCOVY.

## IMPORTANT FACTS

This is only a brief summary of important information about DESCOVY® and does not replace talking to your healthcare provider about your condition and your treatment.

## POSSIBLE SIDE EFFECTS OF DESCOVY

### DESCOVY can cause serious side effects, including:

- Those in the “Most Important Information About DESCOVY” section.
- Changes in your immune system.
- New or worse kidney problems, including kidney failure.
- Too much lactic acid in your blood (lactic acidosis), which is a serious but rare medical emergency that can lead to death. Tell your healthcare provider right away if you get these symptoms: weakness or being more tired than usual, unusual muscle pain, being short of breath or fast breathing, stomach pain with nausea and vomiting, cold or blue hands and feet, feel dizzy or lightheaded, or a fast or abnormal heartbeat.
- Severe liver problems, which in rare cases can lead to death. Tell your healthcare provider right away if you get these symptoms: skin or the white part of your eyes turns yellow, dark “tea-colored” urine, light-colored stools, loss of appetite for several days or longer, nausea, or stomach-area pain.

**The most common side effect of DESCOVY** is nausea.

These are not all the possible side effects of DESCOVY. Tell your healthcare provider right away if you have any new symptoms while taking DESCOVY.

**Your healthcare provider will need to do tests to monitor your health before and during treatment with DESCOVY.**

## HOW TO TAKE DESCOVY

- DESCOVY is a one pill, once a day HIV-1 medicine that is taken with other HIV-1 medicines.
- Take DESCOVY with or without food.

## GET MORE INFORMATION

- This is only a brief summary of important information about DESCOVY. Talk to your healthcare provider or pharmacist to learn more.
- Go to DESCOVY.com or call 1-800-GILEAD-5.
- If you need help paying for your medicine, visit DESCOVY.com for program information.